


## Surgery to Repair Abdominal Wall Hernia

### Information for Patients

#### What is a Hernia?

A Hernia is an out-pouching of intra-abdominal contents through a defect, or hole in the muscle. The abdominal muscles and bones of the pelvis prevent abdominal contents from drooping out under normal circumstances. The skin and fat below the skin do not provide any strength and stretch if there is a hernia present.



Hernias can occur anywhere in the abdomen, but most commonly occur at weak points where a small tear can start and enlarge over time. The common weak points are:

- i) Groin Hernia – These are the commonest and include inguinal and femoral (they are discussed in a separate sheet)
- ii) Umbilical – Hernias around the belly button are common and often present as an ‘outie’ type umbilicus
- iii) Incisional – Scar tissue after previous surgery is weaker than the surrounding muscle and can tear to form a hernia
- iv) Epigastric – Some people get hernias in the region between the belly button and the breastbone.
- v) Lumbar – These occur around the side, often after kidney surgery

#### Predisposing factors

It is not always possible to identify the time when a hernia started, although sometimes that is the case. Hernias are often associated with either weak muscles or raised abdominal pressure (or both).

Frequent features are:

- A history of heavy lifting
- Male Sex
- Obesity
- Chronic cough, constipation, or urinary retention
- Previous surgery
- Divarication of the rectus muscles

#### Symptoms of hernia

Often there are no symptoms of hernia. When present, the common symptoms are:

- A lump that comes and goes and is more prominent on standing or straining.
- Pain, discomfort or ‘dragging’ sensation at the site.
- Incarceration or strangulation (see below).

#### Complications of Hernias

Hernias do not repair themselves. Over time, hernias tend to get bigger as the defect in the muscle stretches to allow more abdominal contents to slip in and out. As time goes on there is a risk that the hernia will develop a complication.

**Incarceration** occurs with a long term (chronic) hernia where the contents come out and stay out and are unable to be pushed back in. This often causes discomfort or mild to moderate pain but not severe pain.

**Bowel Obstruction** occurs when the bowel within the hernia kinks off and blocks.

**Strangulation** is where the contents come out and cannot be pushed back in, and where the defect causes such pressure on the blood vessels that the contents are starved of blood supply. This causes swelling, and severe pain. This is an emergency and warrants prompt attention at the hospital.

#### Who needs a hernia repair?

Patients with hernias, who are fit enough to undergo surgery should have hernias repaired before complications occur. Patients with strangulated hernias require urgent repair.

#### What tests are done?

Most commonly hernias are diagnosed clinically by your doctor. Sometimes ultrasound and/or CT scans can help rule out other causes of symptoms and diagnose the hernia. Tests to ensure safety of anaesthesia are performed according to age group and risk factors.

#### How is the hernia repaired?

The principle of hernia repair is to reduce the contents back into the abdominal cavity and close the muscle. There are various techniques for closing the muscle and Dr Crawford will discuss the options and make recommendations in your instance. In

general, small defects can be closed with a simple stitch technique, whereas larger holes require both stitching and mesh. Very large defects might require pre-operative Botox (to relax the muscles) or releasing incisions in the muscles at the time of surgery.

**Can it be repaired with laparoscopic (keyhole) surgery?**

Yes, this is the most common method of repair, although it is not suitable for everyone. Most non-groin hernias are repaired with a laparoscopic suture and mesh technique.

**Divarication of the Rectus/diastasis recti**

Here the 'six pack' muscles separate above the umbilicus with stretched out tissue. This is sutured closed at the time of the hernia repair.

**The post-operative course**

The post operative course is different for each person. Most patients go home a day or 2 after surgery. Abdominal wall hernia patients are usually placed in a 'binder' (an elasticated corset) that they wear for around 3 weeks.

Skin sutures are dissolving and buried, dressings can stay on for 7 days.

A follow-up appointment is made for 3 weeks after surgery. You may eat and drink normally and walk around straight away. **Heavy lifting and vigorous sports should be avoided for 6 weeks after the surgery.**

**Serious complications after hernia surgery are rare.**

**Complications of hernia surgery include (but not limited to):**

- *Hernia recurrence.* No matter what method of repair is used there is a small lifetime risk of the hernia recurring. Unfortunately, no repair is as good as the original.

- *Bowel injury.* A rare complication of surgery where the bowel is injured during the removal of old scar tissue.
- *Mesh infection.* A serious complication that could lead to further surgery and long recovery
- *Wound infection.*
- *Clots.*
- *Allergic reactions.*
- *Heart troubles.*

**FAQs**

***Will I be able to go to the gym / lift heavy objects again?***

Yes, while there will always be a risk of another hernia, the risks are small after a mesh hernia repair. Dr Crawford will inform you when you are ready to go back to these activities (usually at least 6 weeks after surgery).

***What should I do if the hernia strangulates before surgery can be scheduled?***

If the hernia suddenly becomes swollen and painful or has a red appearance, then you should lie down and try to gently push it back. If this works you should contact your doctor as soon as possible. If this does not work within 30 minutes, then you should go immediately to hospital.

***Does the mesh stay in permanently?***

Yes. In general, the mesh is made of a polypropylene or polyester material that blends with your muscle and promotes scar tissue. It reinforces your muscle and is intentionally not dissolvable. It is very rare for the mesh to need to be removed, but if it were to get infected it might be removed with another operation.

Visit my website or YouTube:  
<https://youtu.be/HbfMhzd9Wby>  
to watch a video

**Please ask Dr Crawford if you have further questions.**

**Dr Michael Crawford**  
**Laparoscopic & Hepatobiliary Surgeon**  
[laparoscopicsurgeon.net.au](http://laparoscopicsurgeon.net.au)

**Phone: (02) 9565 4854**

Suite 314      Fax: (02) 9158 8620      Suite 1.17  
RPA Medical Centre      The Mater Clinic  
100 Carillon Ave      25 Rocklands Rd  
Newtown 2042      Wollstonecraft 2065